

Conditional implementation of a quantum phase gate with distant atoms trapped in different optical cavities

XuBo Zou, and W. Mathis
*Electromagnetic Theory Group at THT,
 Department of Electrical Engineering,
 University of Hannover, Germany*

We propose a scheme for conditional implementation of a quantum phase gate by using distant atoms trapped in different optical cavities. Instead of direct interaction between atoms, the present scheme makes use of quantum interference of polarized photons decaying from the optical cavities to conditionally create the desired quantum phase gate between two distant atoms. The proposed scheme only needs linear optical elements and a two-fold coincidence detection, and are insensitive to the quantum noise. The scheme can be directly used to prepare any quantum state of many distant atoms.

PACS numbers: 03.65.Ud, 42.50.-p

The manipulation and control of quantum entanglement between many distant atoms is an important challenge for implementation of quantum communication and computation[1]. Recently, several physical systems have been suggested as possible candidates for engineering quantum entanglement: cavity QED systems [2], trapped ion systems [3], quantum dot systems [4] and Josephson-junction device systems[5]. Among them, cavity QED systems are paid more attention. This is due to the fact that cold and localized atoms are well suitable for storing quantum information in long-lived internal states, and the photons are natural source for fast and reliable transport of quantum information over long distance. A number of scheme have been proposed for manipulating quantum entanglement between atoms and cavity fields[6, 7]. In particular, entangled states of two or three particles have been demonstrated experimentally in high-Q cavities [8, 9]. Schemes of this type are based on the controlling of direct or indirect interaction between the atoms, which are intended to be entangled. Since most of these schemes require a high-Q cavity field, decoherence caused by cavity decay can be neglected.

Conditional measurements offer another possible way to engineer quantum entanglement and implement quantum information processing. In Ref[10], Cabrillo et al proposed a scheme to entangle quantum states of spatial widely separated atoms by weak laser pulses and the detection of the subsequent spontaneous emission. Further, several schemes have also been proposed for generating the EPR state between two distant atoms trapped in different optical cavities[11, 12]. In Ref[13], schemes were also proposed for generating quantum entanglement of many distant atoms. However these schemes are not easily extended to generate any quantum entanglement. In order to generate any states and implement quantum algorithm on demand, it is necessary to propose a scheme to implement quantum logic gates. The universality of the set of the controlled-NOT gate (or controlled-phase gate) and single-qubit gates implies that we can create any states and implement any quantum algorithm by constructing a quantum circuit using these gates. In Ref[14], Protsenko et al proposed a scheme for conditional implementation of a quantum controlled-NOT gate between two distant atoms by the detection of single scattered photon. A disadvantage of this schemes is to require single-photon detector to distinguish zero photon, one photon and more than one photon. The inefficiency of photon detector influences the fidelity of the quantum operations. Another disadvantage of the scheme is to require that the laser pulses have to be sufficiently weak to ensure that the probability of exciting both atoms is much smaller the probability to excite only one atom. In such case, it will be a time-consuming task to detect single scattered photon. In this paper, we propose an alternative scheme to implement conditional quantum phase gate with two distant atoms trapped in different optical cavities with the two-fold coincidence detection. The protocol has the following favorable features. (1) The scheme is insensitive to the imperfection of the photon detectors, i.e. the scheme does not require distinguishing between zero, one and two photons. (2) The scheme is insensitive to the phase accumulated by the photons on their way from the cavities to the place where they are detected. (3)The scheme does not the weakly driven laser pulses as both atoms are excited simultaneously. (4) The scheme can be used to prepare any quantum entanglement of many distant atoms. The quantum entanglement between many distant sites is an essential element of quantum network construction.

The schematic representation of our scheme for conditional quantum phase gate is shown in Fig.1, which consists of two identical atoms 1 and 2 confined separately in two optical cavities 1 and 2, respectively. The photons leaking out from the cavities 1 and 2 are interfered at a polarization beam splitter (PBS), with the outputs detected by four single-photon detectors after two quarter wave plates (QWP) and two polarization beam splitters. For the photons decaying from the cavity 1, a quarter wave plate is inserted before the first polarization beam splitter. As shown in Fig.2, each atom (index $j = 1, 2$) has four Zeeman sublevels $|g_H\rangle_j$, $|g_V\rangle_j$, $|s_H\rangle_j$, $|s_V\rangle_j$ and two excited state $|e_H\rangle_j$, $|e_V\rangle_j$. The lifetime of the atomic levels $|g_H\rangle_j$, $|g_V\rangle_j$, $|s_H\rangle_j$, $|s_V\rangle_j$ is assumed to be comparatively long so that

spontaneous decay of these states can be neglected. We encode the ground states $|g_H\rangle_j$ and $|g_V\rangle_j$ as logic zero and one states, i.e. $|g_H\rangle_j = |0\rangle_j$ and $|g_V\rangle_j = |1\rangle_j$. Such an atomic level structure has been proposed to implement quantum computation in a single cavity [15]. The transitions $|g_H\rangle_j \rightleftharpoons |e_H\rangle_j$ and $|g_V\rangle_j \rightleftharpoons |e_V\rangle_j$ are coupled to two degenerate cavity modes a_{jH} and a_{jV} with different polarization H and V , respectively. The transitions $|e_H\rangle_j \rightleftharpoons |s_H\rangle_j$ and $|e_V\rangle_j \rightleftharpoons |s_V\rangle_j$ are driven by two classical fields, respectively. We assume that the classical laser fields and the cavity fields are detuned from their respective transitions by the same amount. In the case of large detuning the excited states can be eliminated adiabatically to obtain the effective interaction Hamiltonian (in the interaction picture)

$$H_j = \Omega(a_{jH}|g_H\rangle_{jj}\langle s_H| + a_{jH}^\dagger|s_H\rangle_{jj}\langle g_H| + a_{jV}|g_V\rangle_{jj}\langle s_V| + a_{jV}^\dagger|s_V\rangle_{jj}\langle g_V|) \quad (1)$$

where a_{jH} and a_{jV} ($a_{jH}^\dagger$ and $a_{jV}^\dagger$) are the annihilation (creation) operators of the H and V polarization modes of the j th cavity. Here we assumed that the effective coupling constants of the atoms coupled with their cavity modes are same, which are described by Ω . In order to investigate the quantum dynamics of the system, it is convenient to follow a quantum trajectory description [16]. The evolution of the system's wave function is governed by a non-Hermitian Hamiltonian

$$H'_j = H_j - i\kappa(a_{jH}^\dagger a_{jH} + a_{jV}^\dagger a_{jV}) \quad (2)$$

as long as no photon decays from the cavity. Here we assume that two optical cavities have the same loss rate κ for the all modes. If a single-photon detector D_j ($j = 1, 2, 3, 4$) detects a photon, the coherent evolution according to H'_j is interrupted by a quantum jump. This corresponds to a quantum jump, which can be formulated with the operators b_j on the joint state vectors of two atom-cavity systems

$$\begin{aligned} b_1 &= \frac{1}{2}(a_{1H} + a_{1V}) + \frac{1}{\sqrt{2}}a_{2V}, \quad b_2 = \frac{1}{2}(a_{1H} + a_{1V}) - \frac{1}{\sqrt{2}}a_{2V} \\ b_3 &= \frac{1}{2}(a_{1H} - a_{1V}) + \frac{1}{\sqrt{2}}a_{2H}, \quad b_4 = -\frac{1}{2}(a_{1H} - a_{1V}) + \frac{1}{\sqrt{2}}a_{2H} \end{aligned} \quad (3)$$

In the following, we analyze the scheme in detail. In order to demonstrate the conditional implementation of quantum phase gate, we assume that atom, trapped in the j -th ($j = 1, 2$) optical cavity, are initially prepared in the state

$$\alpha_j|g_H\rangle_j + \beta_j|g_V\rangle_j, \quad (4)$$

and both polarization modes of optical cavity are prepared in the vacuum states $|0, 0\rangle_j$, where $|m, n\rangle_j$ denotes m photons in polarization mode and n photons in the polarization mode. Now we switch on the Hamiltonian(1) in each atom-cavity system for a time τ . If no photon is emitted from the cavity, the j th atom-cavity system is governed by the interaction H'_j . In this case the atom-cavity state evolves to the entangled state

$$|\Psi\rangle_j = \alpha_j[a|g_H\rangle_j|0, 0\rangle_j - ib|s_H\rangle_j|1, 0\rangle_j] + \beta_j[a|g_V\rangle_j|0, 0\rangle_j - ib|s_V\rangle_j|0, 1\rangle_j] \quad (5)$$

with

$$\begin{aligned} a &= \frac{\cos(\Omega_\kappa\tau) - \frac{\sin(\Omega_\kappa\tau)\kappa}{2\Omega_\kappa}}{\sqrt{[\cos(\Omega_\kappa\tau) - \frac{\sin(\Omega_\kappa\tau)\kappa}{2\Omega_\kappa}]^2 + \frac{\sin^2(\Omega_\kappa\tau)\Omega^2}{\Omega_\kappa^2}}} \\ b &= \frac{\sin(\Omega_\kappa\tau)\Omega}{\Omega_\kappa\sqrt{[\cos(\Omega_\kappa\tau) - \frac{\sin(\Omega_\kappa\tau)\kappa}{2\Omega_\kappa}]^2 + \frac{\sin^2(\Omega_\kappa\tau)\Omega^2}{\Omega_\kappa^2}}} \\ \Omega_\kappa &= \sqrt{\Omega^2 - \kappa^2/4} \end{aligned} \quad (6)$$

The probability that no photon is emitted during this evolution becomes

$$P_{single} = e^{-\kappa\tau} \left\{ [\cos(\Omega_\kappa\tau) - \frac{\sin(\Omega_\kappa\tau)\kappa}{2\Omega_\kappa}]^2 + \frac{\sin^2(\Omega_\kappa\tau)\Omega^2}{\Omega_\kappa^2} \right\} \quad (7)$$

We assume that the interaction Hamiltonian (1) is applied to each atom-cavity system simultaneously, so that the atom-cavity states $|\Psi\rangle_j$ ends at the same time. This implements the first step of the protocol. The probability that this stage is a success is the probability that no photon decays from either atom-cavity system during the preparation. This quantity is given by $P_{suc} = P_{single}^2$.

Next we consider the second step of the scheme, in which we make a photon number measurement with four single-photon detectors D_j ($j = 1, 2, 3, 4$) on the output modes of the setup. In this step, the joint state of two atom-cavity systems becomes prepared in the form

$$|\Phi(0)\rangle = |\Psi\rangle_1 \otimes |\Psi\rangle_2 \quad (8)$$

where the state $|\Psi\rangle_j$ is given by Eq.(5). We assume that photons are detected at the time t . This assumption is posed to calculate the system's time evolution during this time interval in a consistent way with the "no-photon-emission-Hamiltonian" (2). The joint state of the total system evolves into

$$|\Phi(t)\rangle = |\Psi(t)\rangle_1 \otimes |\Psi(t)\rangle_2 \quad (9)$$

with

$$|\Psi(t)\rangle_j = \frac{1}{a^2 + b^2 e^{-2\kappa t}} \{ \alpha_j [a|g_H\rangle_j |0, 0\rangle_j - ibe^{-\kappa t - i\varphi_j} |s_H\rangle_j |1, 0\rangle_j] + \beta_j [a|g_V\rangle_j |0, 0\rangle_j - ibe^{-\kappa t - i\varphi_j} |s_V\rangle_j |0, 1\rangle_j] \} \quad (10)$$

where the phases $\varphi_j = kL_j$, k is the wave number and L_j are the optical lengths which photons travel through the optical system towards the photon detectors. The detection of one photon with the detector D_j corresponds to a quantum jump, which can be formulated with the operator b_j on the joint state $|\Phi(t)\rangle$. If D_1 and D_3 detect one photon at nearly the same time and D_2 and D_4 do not detect any photon, or vice, the state of the total system is projected into

$$\alpha_1 \alpha_2 |s_H\rangle_1 |s_H\rangle_2 + \beta_1 \alpha_2 |s_V\rangle_1 |s_H\rangle_2 + \alpha_1 \beta_2 |s_H\rangle_1 |s_V\rangle_2 - \beta_1 \beta_2 |s_V\rangle_1 |s_V\rangle_2 \quad (11)$$

If D_1 and D_4 detect one photon at nearly the same time and D_2 and D_3 do not detect any photon during that time interval, or vice, the state of the total system becomes projected into

$$\alpha_1 \alpha_2 |s_H\rangle_1 |s_H\rangle_2 + \beta_1 \alpha_2 |s_V\rangle_1 |s_H\rangle_2 - \alpha_1 \beta_2 |s_H\rangle_1 |s_V\rangle_2 + \beta_1 \beta_2 |s_V\rangle_1 |s_V\rangle_2 \quad (12)$$

where we have neglected the multiplicative factor $e^{i(\varphi_1 + \varphi_2)}$ in Eq.(11) and (12). This result demonstrates that phase accumulated by the photons have no effect on the conditional implementation of the quantum operation. Based on the result of measurement, we can apply fast Raman transitions to individually manipulate the atoms 1 and 2, and map the states Eq.(11) or Eq.(12) to the state

$$\alpha_1 \alpha_2 |g_H\rangle_1 |g_H\rangle_2 + \beta_1 \alpha_2 |g_V\rangle_1 |g_H\rangle_2 + \alpha_1 \beta_2 |g_H\rangle_1 |g_V\rangle_2 - \beta_1 \beta_2 |g_V\rangle_1 |g_V\rangle_2 \quad (13)$$

which demonstrates the conditional implementation of quantum controlled phase gate

$$\begin{aligned} |g_H\rangle |g_H\rangle &\rightarrow |g_H\rangle |g_H\rangle, & |g_H\rangle |g_V\rangle &\rightarrow |g_H\rangle |g_V\rangle \\ |g_V\rangle |g_H\rangle &\rightarrow |g_V\rangle |g_H\rangle, & |g_V\rangle |g_V\rangle &\rightarrow -|g_V\rangle |g_V\rangle \end{aligned} \quad (14)$$

The success probability of the scheme is $P_s = b^4(1 - e^{-2\kappa t})^2/2$. We now give a brief discussion on the influence of practical noise on the scheme. In the present scheme, the lifetime of the atomic ground levels is assumed to be comparatively long so that spontaneous decay of these states can be neglected. The dominate noise of the scheme is the photon loss, which includes the contribution from channel attenuation, and the inefficiency of the single-photon detectors. All these kinds of noise can be considered by an overall photon loss probability η [11]. It is noticed that the present scheme is based on the two-photon coincidence detection. If one photon is lost, a click from each of the detectors is never recorded. In this case, the scheme fails to generate the expected quantum operation. Therefore the photon loss only decreases the success probability P_s by a factor of $(1 - \eta)^2$, but have no influence on the fidelity of the expected operation.

In summary, we have proposed a scheme to implement the conditional quantum phase gate for two distant atoms trapped in different optical cavities by using interference of polarization photons. Compared with previous scheme[14], the scheme is insensitive to the imperfection of the photon detectors. The detection efficiency does not affect the fidelity of the conditional quantum operation and photon counting detectors are not needed. Low detection efficiency lowers only the success probability. The most possible application of the present scheme is to create quantum

entanglement between many distant atoms trapped in different optical cavities. Recently, researchers are turning their interest towards characterizing and generating multipartite entanglement, and using it for more general and useful applications[17]. Although several schemes have been proposed for quantum entanglement of many distant atoms[13], these schemes are not easily extended to generate any quantum entanglement. By employing the quantum logic networks proposed in Ref[18], the present scheme can be directly used to conditionally generate any quantum state of many distant atoms.

-
- [1] P. Shor in Proceeding of the 35th Annual symposium on the foundation of Computer Science (IEEE Comp. Soc. Press, Los Alomitos, CA, 1994) 124-134; Lov K. Grover Phys. Rev. Lett. 80, 4329 (1998); A. Einstein, B. Podolsky and N. Rosen, Phys. Rev. 47, 777 (1935); J. S. Bell, Physics (Long Island City, N.Y.) 1, 195 (1965); A. K. Ekert, Phys. Rev. Lett. 67, 661 (1991); C. H. Bennett and S.J. Wiesner, Phys. Rev. Lett. 69, 2881 (1992); C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. Wootters, Phys. Rev. Lett. 70, 1895 (1993)
 - [2] Q. A. Turchette, C. J. Hood, W. Lange, H. Mabuchi and H. J. Kimble Phys. Rev. Lett. 75(1995)4710
 - [3] C. Monroe, D. M. Meekhof, B. E. King, W.M. Itano and D. J. Wineland, Phys. Rev. Lett. 75(1995)4714; J. I. Cirac and P. Zoller, Phys. Rev. Lett. 74(1995)4091
 - [4] D. Loss and D. P. Divincenzo, Phys. Rev. A. 57(1998)120
 - [5] Y. Makhlin and G. Schon, nature 398(1999)305
 - [6] J. M. Raimond et al., Rev. Mod. Phys. 73, 565 (2001)
 - [7] J. I. Cirac and P. Zoller, Phys. Rev. A 50, R2799 (1994); C. C. Gerry, Phys. Rev. A 53, 2857 (1996); W. Lange and H. J. Kimble, Phys. Rev. A 61, 063817 (2000)
 - [8] E. Hagley, X. Maitre, G. Nogues, C. Wunderlich, M. Brune, J. M. Raimond, and S. Hroche, Phys. Rev. Lett. 79, 1 (1997).
 - [9] A. Rauschenbeutel et al, Science.288, 2024 (2000).
 - [10] C. Cabrillo, J. I. Cirac, P. Garcia-Fernandez, and P. Zoller, Phys. Rev. A 59, 1025 (1999)
 - [11] S. Bose, P. L. Knight, M. B. Plenio and V. Vedral, Phys. Rev. Lett. 83, 5158 (1999)
 - [12] X. L. Feng et al Phy. Rev. Lett. 90, 217902 (2003); L. M. Duan and H. J. Kimble, Phy. Rev. Lett. 90, 253601 (2003); D. E. Browne, M. B. Plenio, and S. F. Huelga, Phys. Rev. Lett. 91, 067901 (2003); C. Simon and W. T. M. Irvine, Phys. Rev. Lett. 91, 110405 (2003)
 - [13] XuBo Zou, K. Pahlke and W. Mathis, Phys. Rev. A 68, 024302 (2003); C. D. Fidio and W. Vogel, J.Opt.B 5, 105 (2003)
 - [14] I. E. Protsenko, G. Reymond, N. Schlosser, P. Grangier, Phys. Rev. A 66, 062306 (2002)
 - [15] T. Pellizzari, S. A. Gardiner, J. I. Cirac, and P. Zoller, Phys. Rev. Lett. 75, 3788(1995)
 - [16] M. B. Plenio and P. L. Knight, Rev. Mod. Phys. 70, 101 (1998).
 - [17] D. M. Greenberger, M. Horne, A. Shimony, and A. Zeilinger, Am. J. Phys. 58, 1131 (1990); W. Duer, G. Vidal, and J.I. Cirac, Phys. Rev. A 62, 062314 (2000)
 - [18] M. Sasura and V. Buzek, Phys. Rev. A 64, 012305 (2001)

Figure Captions

Figure 1. The schematic setup for conditional quantum phase gate of two distant atoms trapped in different optical cavities, which includes three polarization beam splitters (PBS), three quarter wave plates (QWP) and four single-photon detectors.

Figure 2. The relevant level structure with ground state $|g_H\rangle$, $|s_H\rangle$, $|g_V\rangle$, $|s_V\rangle$ and excited states $|e_H\rangle$, $|e_V\rangle$. The transitions $|e_H\rangle \rightarrow |s_H\rangle$ and $|e_V\rangle \rightarrow |s_V\rangle$ are coupled by the classical lasers and the transitions $|e_H\rangle \rightarrow |g_H\rangle$ and $|e_V\rangle \rightarrow |g_V\rangle$ are coupled to two degenerate cavity modes a_H and a_V with different polarization H and V .